arXiv:1005.4256v1 [math.CO] 24 May 2010

Bijective Proofs of Gould's and Rothe's Identities

Victor J. W. Guo[1]

Department of Mathematics, East China Normal University, Shanghai 200062, People's Republic of China
`jwguo@math.ecnu.edu.cn`, `http://math.ecnu.edu.cn/~jwguo`

AMS Subject Classifications: 05A19

Abstract. We first give a bijective proof of Gould's identity in the model of binary words. Then we deduce Rothe's identity from Gould's identity again by a bijection, which also leads to a double-sum extension of the q-Chu-Vandermonde formula.

Keywords: Rothe's identity, Gould's identity, binary words, bijection

1 Introduction

There are two convolution formulas due to Rothe [10]:

$$\sum_{k=0}^{n}\frac{xy}{(x-kz)(y-(n-k)z)}\binom{x-kz}{k}\binom{y-(n-k)z}{n-k}=\frac{x+y}{x+y-nz}\binom{x+y-nz}{n}, \tag{1}$$

$$\sum_{k=0}^{n}\frac{x}{x-kz}\binom{x-kz}{k}\binom{y+kz}{n-k}=\binom{x+y}{n}, \tag{2}$$

which are famous in the literature. For example, Chu [3] used (1) and (2) to compute some determinants involving binomial coefficients. For some generalizations of (1) and (2), we refer the reader to [11, 13, 14] and references therein.

Some proofs of (1) and (2) can be found in [6, 9, 12]. It is not difficult to see that (1) can be deduced from (2). Blackwell and Dubins [2] have given a combinatorial proof of Rothe's identity (1), which can also be proved in the model of lattice paths (using [8, p. 9] or [7, (1.1)]).

Gould [4, 5] reproved (1) and (2) and also obtained the following interesting identity:

$$\sum_{k=0}^{n}\binom{x-kz}{k}\binom{y+kz}{n-k}=\sum_{k=0}^{n}\binom{x+\epsilon-kz}{k}\binom{y-\epsilon+kz}{n-k}. \tag{3}$$

The main purpose of this paper is to give bijective proofs of Rothe's identity (2) and Gould's identity (3) in the model of binary words. As a conclusion, a double-sum extension of the q-Chu-Vandermonde formula is also presented.

---

[1] The author was partially supported by a Junior Research Fellowship at the Erwin Schrödinger International Institute for Mathematical Physics in Vienna.

2 Proof of (3)

It is not difficult to see that Gould's identity (3) is equivalent to

$$\sum_{k=0}^{n} \binom{p-km}{k}\binom{q+km}{n-k} = \sum_{k=0}^{n} \binom{p+1-km}{k}\binom{q-1+km}{n-k} \quad \text{for } p,q,m,n \in \mathbb{N}. \tag{4}$$

We will prove that (4) holds for all integers $p \geq mn$ and $q \geq 1$ (and therefore for all real and complex numbers).

Let $\Gamma = \{a, b\}$ denote an alphabet with a grading $||a|| = 1$ and $||b|| = m+1$. For a word $w = w_1 \cdots w_n \in \Gamma^*$, its *length* n is denoted by $|w|$ and its *weight* by $||w|| = ||w_1|| + \cdots + ||w_n||$. Let $|w|_b = (||w|| - |w|)/m$ be the number of b's appearing in w, and let

$$\Gamma_{p,k} := \{w \in \Gamma^* \colon ||w|| = p \text{ and } |w|_b = k\} \subseteq \Gamma^{p-km}.$$

It is easy to see that $\#\Gamma_{p,k} = \binom{p-km}{k}$. Furthermore, let

$$\Gamma_{p,k}^{(r)} := \{w \in \Gamma_{p,k} \colon w \text{ has a prefix of weight } r\}.$$

For $p, q \geq mn$, an obvious bijection (by factorization)

$$\Gamma_{p+q,n}^{(p)} \longleftrightarrow \biguplus_{k} \Gamma_{p,k} \times \Gamma_{q,n-k}$$

leads to

$$\#\Gamma_{p+q,n}^{(p)} = \sum_{k} \binom{p-km}{k}\binom{q-(n-k)m}{n-k}.$$

Thus, the identity (4) is equivalent to

$$\#\Gamma_{p+q+mn,n}^{(p)} = \#\Gamma_{p+q+mn,n}^{(p+1)}. \tag{5}$$

We need the following simple observation.

Lemma 1 *Let $u, v \in \Gamma^*$ with $||u||, ||v|| \geq mn+1$, where $n = |u \cdot v|_b$. Then there exist nonempty prefixes x of u and y of v such that $||x|| = ||y||$.*

Proof. Suppose that $||u|| = mn + r$ and $||v|| = mn + s$ with $r, s \geq 1$. Then the total number of nonempty prefixes of u and v is $|u| + |v| = ||u|| + ||v|| - m|u \cdot v|_b = mn + r + s$. On the other hand, each prefix of u or v has weight $\leq \max\{||u||, ||v||\}$. Hence u and v must have some nonempty prefixes of the same weight. ∎

Now we can prove (5) by the following theorem.

Theorem 2 *For $p \geq mn$ and $q \geq 1$, there is a bijection between $\Gamma_{p+q+mn,n}^{(p)}$ and $\Gamma_{p+q+mn,n}^{(p+1)}$.*

Proof. Take any $w = u \cdot v \in \Gamma_{p+q+mn,n}^{(p)}$, where $||u|| = p$ and $||v|| = q + mn$. Applying Lemma 1 to v and the reverse of $u \cdot a$, we see that u has a suffix x (possibly empty), i.e., $u = u' \cdot x$, and v has a prefix y, i.e., $v = y \cdot v'$, such that $||x|| = ||y|| - 1$. Then $u' \cdot \overline{y} \cdot \overline{x} \cdot v' \in \Gamma_{p+q+mn,n}^{(p+1)}$, where $\overline{x}$ and $\overline{y}$ are respectively the reverses of x and y. By selecting x and y with minimal length, we obtain a bijection. ∎

3 Proof of (2)

Let us now consider $\Gamma_{p+q+mn,n}$ with $p \geq mn$ and $q \geq 1$. For each $w \in \Gamma_{p+q+mn,n}$, let $w = u \cdot v$ denote the unique factorization with $||u|| \geq p$ but as small as possible. There are two possibilities:

- If $||u|| = p$, then $w \in \Gamma^{(p)}_{p+q+mn,n}$ and all these words have been counted above.
- If $||u|| = p + j$ for some $1 \leq j \leq m$, then the last letter of u must be a b. Namely, $u = u' \cdot b$ for some $u' \in \Gamma_{p+j-m-1,k-1}$. The corresponding v belongs to $\Gamma_{q+mn-j,n-k}$. It is easy to see that the mapping $w \mapsto (u', v)$ may be inverted.

Hence there is a bijection

$$\Gamma_{p+q+mn,n} \longleftrightarrow \Gamma^{(p)}_{p+q+mn,n} \biguplus_{j=1}^{m} \biguplus_{k=1}^{n} \Gamma_{p+j-m-1,k-1} \times \Gamma_{q+mn-j,n-k}, \tag{6}$$

which means that

$$\sum_{k=0}^{n} \left(\binom{p-km}{k}\binom{q+km}{n-k} + \sum_{j=1}^{m} \binom{p-km+j-1}{k-1}\binom{q+km-j}{n-k} \right) = \binom{p+q}{n}. \tag{7}$$

However, by (4), for $1 \leq j \leq m$, we have

$$\sum_{k=0}^{n} \binom{p-km+j-1}{k-1}\binom{q+km-j}{n-k} = \sum_{k=0}^{n} \binom{p-km-1}{k-1}\binom{q+km}{n-k}. \tag{8}$$

Combining (7) and (8) yields

$$\sum_{k=0}^{n} \left(\binom{p-km}{k} + m\binom{p-km-1}{k-1} \right) \binom{q+km}{n-k} = \binom{p+q}{n},$$

or

$$\sum_{k=0}^{n} \frac{p}{p-km}\binom{p-km}{k}\binom{q+km}{n-k} = \binom{p+q}{n},$$

which is Rothe's identity (2).

4 A new extension of the q-Chu-Vandermonde formula

In this section we give a q-analogue of (7). Recall that *q-binomial coefficient* $\begin{bmatrix} x \\ k \end{bmatrix}$ is defined as $\prod_{i=1}^{k}(1-q^{x-i+1})/(1-q^i)$ if $k \geq 0$ and 0 otherwise. By [1, Theorem 3.6], we have

$$\sum_{w \in \Gamma_{p,k}} q^{\mathrm{inv}(w)} = \begin{bmatrix} p-km \\ k \end{bmatrix} \quad (p \geq km), \tag{9}$$

where $\mathrm{inv}(w)$ denotes the number of inversions of w. Taking $\mathrm{inv}(w)$ into account and using (9), the bijection (6) (replacing p and q by x and y, respectively) further implies that

$$\sum_{k=0}^{n} q^{k(km+k+y-n)} \left(\begin{bmatrix} x-km \\ k \end{bmatrix}\begin{bmatrix} y+km \\ n-k \end{bmatrix} + \sum_{j=1}^{m} \begin{bmatrix} x-km+j-1 \\ k-1 \end{bmatrix}\begin{bmatrix} y+km-j \\ n-k \end{bmatrix} q^{-kj} \right) = \begin{bmatrix} x+y \\ n \end{bmatrix}, \tag{10}$$

which reduces to the q-Chu-Vandermonde formula if $m = 0$, and reduces to

$$\sum_{k=0}^{n} q^{k(2k+y-n)} \left(\begin{bmatrix} x-k \\ k \end{bmatrix} \begin{bmatrix} y+k \\ n-k \end{bmatrix} + \begin{bmatrix} x-k \\ k-1 \end{bmatrix} \begin{bmatrix} y+k-1 \\ n-k \end{bmatrix} q^{-k} \right) = \begin{bmatrix} x+y \\ n \end{bmatrix}$$

if $m = 1$.

However, our bijection in Theorem 2 does not lead to the corresponding q-analogue of (8), and we cannot simplify (10) to obtain a q-analogue of Rothe's identity as before.

Acknowledgments. The author is indebted to one of the referees for his detailed constructive comments and suggestions, which enable the author to shorten this note more than half. Especially, the present form of Lemma 1 is his and it unifies two previous lemmas in this paper.

References

[1] G. E. Andrews, The Theory of Partitions, Cambridge University Press, Cambridge, 1998.

[2] D. Blackwell and L. Dubins, An elementary proof of an identity of Gould's, Bol. Soc. Mat. Mexicana **11** (1966), 108–110.

[3] W. Chu, Binomial convolutions and determinant identities, Discrete Math. **204** (1999), 129–153.

[4] H. W. Gould, Some generalization of Vandermonde's convolution, Amer. Math. Monthly **63** (1956), 84–91.

[5] H. W. Gould, Final analysis of Vandermonde's convolution. Amer. Math. Monthly **64** (1957), 409–415.

[6] I-C. Huang, Inverse relations and Schauder bases, J. Combin. Theory, Ser. A **97** (2002), 203–224.

[7] C. Krattenthaler, Counting lattice paths with a linear boundary, I, Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II **198** (1989), 87–107.

[8] S. G. Mohanty, Lattice Path Counting and Applications, Academic Press, New York, 1979.

[9] G. N. Raney, Functional composition patterns and power series reversion, Trans. Amer. Math. Soc. **94** (1960), 441–451.

[10] H. A. Rothe, Formulae de serierum reversione demonstratio universalis signis localibus combinatorio-analyticorum vicariis exhibita, Leipzig, 1793.

[11] M. Schlosser, Abel-Rothe type generalizations of Jacobi's triple product identity, in: Theory and Applications of Special Functions, Dev. Math., 13, Springer, New York, 2005, pp. 383–400.

[12] R. Sprugnoli, Riordan arrays and the Abel-Gould identity, Discrete Math. **142** (1995), 213–233.

[13] V. Strehl, Identities of Rothe-Abel-Schläfli-Hurwitz-type, Discrete Math. **99** (1992), 321–340.

[14] J. Zeng, Multinomial convolution polynomials, Discrete Math. **160** (1996), 219–228.